



E699

JACC March 12, 2013

Volume 61, Issue 10



## Heart Failure

### MILRINONE PLASMA CONCENTRATIONS ACROSS A SPECTRUM OF RENAL FUNCTION IN STAGE D HEART FAILURE PATIENTS

Poster Contributions

Poster Sessions, Expo North

Sunday, March 10, 2013, 9:45 a.m.-10:30 a.m.

Session Title: Heart Failure: Pharmacologic Therapy

Abstract Category: 17. Heart Failure: Therapy

Presentation Number: 1223-300

Authors: *Zac Cox, Marion W. Calcutt, Thomas B. Morrison, Scott Akers, Mary Beth Davis, Daniel J. Lenihan, Vanderbilt University Medical Center, Nashville, TN, USA, Lipscomb University College of Pharmacy, Nashville, TN, USA*

**Background:** Milrinone exhibits a sigmoidal relationship between plasma concentrations and the percent increase of cardiac index, with no additional hemodynamic benefit above 500 ng/ml. Milrinone may accumulate in renal dysfunction and increase arrhythmic risk in a concentration dependent manner. We hypothesized that heart failure (HF) patients with renal dysfunction would have elevated steady-state milrinone concentrations.

**Methods:** 29 patients on continuous milrinone infusion at the time of plasma collection were enrolled in the Vanderbilt Heart Registry. Creatinine clearance (CrCl) was used to stratify patients into four groups: Group 1 (CrCl >60ml/min), Group 2 (CrCl 60-30 ml/min), Group 3 (CrCl <30 ml/min), and Group 4 (hemodialysis). Plasma milrinone concentration was determined by liquid chromatography-mass spectrometry. Hemodynamic and arrhythmic effects were compared to a pre-milrinone baseline.

**Results:** All patients (76% male; 39% ischemic cardiomyopathy) had stage D HF and an implanted cardiac defibrillator (ICD). There was no difference in the mean infusion rates ( $0.391 \pm 0.08$  mcg/kg/min) ( $p=0.14$ ) between groups. Of 8 post-milrinone ventricular tachycardia episodes requiring defibrillation, 6 occurred in Group 4 patients.

**Conclusions:** Milrinone concentrations are markedly elevated in renal dysfunction. Quantification of milrinone concentrations may identify those at highest risk of ventricular arrhythmias.

	Group 1 (n=14)	Group 2 (n=10)	Group 3 (n=3)	Group 4 (n=2)
Milrinone (ng/ml)	451 <sup>a</sup> ± 243	591 ± 293	1,575 ± 962	6,252 ± 4,409 <sup>b</sup>
CI increase <sup>a</sup>	24% ± 25%	44% ± 45%	50% ± 14%	15% ± 35%

Data as mean ± SD

<sup>a</sup> Cardiac Index

<sup>b</sup>  $p<0.05$  vs all groups